



ORAL CHALLENGE PROCEDURE

Your oral challenge to _____ is scheduled for _____ at _____ AM/PM.

What is an oral challenge?

This test may help confirm or disprove that you have an allergy to a specific medication or food.

An oral challenge can only be performed if you/your child are in good health. If there is any significant rash, fever, cold symptoms, or worsening asthma, please call to reschedule. The patient should be symptom-free for *at least one week*. If in doubt, please call and discuss this with a nurse.

If the patient is <18 yrs. old, a **parent/legal guardian should be present** for the entire Oral Challenge.

If this is not feasible, please complete the “*Consent to treat a minor without a parents/legal guardian present for an oral challenge.*” If this form is not presented at the appointment, we will ask you to reschedule.

What happens during a challenge?

You/your child will ingest increasing amounts of the challenge item (i.e., food or drug) and be observed for a set time between each dose and a more extended period after the last dose.

Are there any long-term effects of an oral challenge?

The test may cause you/your child to develop hives/rash, wheeze, chest tightness, shortness of breath, abdominal pain, or nausea/vomiting. Potentially life-threatening reactions could occur. There has been one documented *death* from an oral challenge. You/your child may be given medication(s) to relieve the symptoms rapidly. There are no known long-term side effects.

How to prepare for your oral challenge?

You/your child **may** be asked to stop antihistamines for 3-10 days before the test (*see chart on next page for details*). If *skin testing* is also being done, please stop antihistamines beforehand.

Allergist preference Stop antihistamine Do not stop antihistamine

- **Please call our office if you have questions about which medications should be held or which may be continued.**
- Do **NOT** stop steroid medications, such as asthma medications, inhalers, nasal steroid sprays, oral steroids (prednisone), or topical steroids.
- Do **NOT** stop other daily non-allergy medications.
- Continue certain reflux medications such as esomeprazole (Nexium), lansoprazole (Prevacid), or omeprazole (Prilosec). *See the chart below* for particular reflux medications to hold.



WHEN TO STOP MEDICATION BEFORE TEST	NAME OF MEDICATION
10 days	Oral antihistamines <ul style="list-style-type: none"> • Cyproheptadine (Periactin) • Hydroxyzine (Atarax, Vistaril)
3-7 days	Nasal spray and/or eye drop antihistamines <ul style="list-style-type: none"> • Azelastine (Astelin, Astepro, Dymista) • Olopatadine (Pataday, Patanase, Patanol) Oral antihistamines (can be in cold/flu/sleep medications) <ul style="list-style-type: none"> • Acrivastine (Semprex-D) • Brompheniramine (in combination products) • Carbinoxamine (Dimetapp, Palgic, Rondec) • Cetirizine (Zyrtec, Wal-Zyr, Allertec) • Chlorpheniramine (Chlor-Trimetron, Triaminic) • Clemastine (Tavist) • Desloratadine (Clarinex) • Diphenhydramine (Benadryl, Nyquil, may end in -PM) • Fexofenadine (Allegra, Allerfex) • Levocetirizine (Xyzal) • Loratadine (Alavert, Allerclear, Claritin) Motion sickness pills: <ul style="list-style-type: none"> • Cyclizine (Marezine, Nausicalm, Valoid) • Meclizine (Antivert, Bonine, Dramamine) Anti-nausea pills: Promethazine (Phenergan)
24 hrs	Anti-reflux medications (which are antihistamines) <ul style="list-style-type: none"> • Cimetidine (Tagamet) • Famotidine (Pepcid)

What to Do on the Day of the Oral Challenge?

- Do **not** use albuterol (Proair, Proventil, Ventolin) or levalbuterol (Xopenex). However, if you/your child have acute asthma symptoms, **please use the rescue inhaler and call the office** to see if rescheduling is necessary.
- Avoid one-piece outfits if skin testing is being done, so arms or back may be accessed.
- Wear comfortable clothing. Consider wearing short-sleeved or sleeveless tops.
- Please avoid eating a large meal at least 2 hours before the visit. You/your child may be permitted to eat about *1 hr after* the last challenge dose.
- You may bring *clear liquids* (Jell-O, fruit juices without pulp, water, iced tea, etc.) to ingest during the challenge. However, breastfeeding may continue as usual.
- Please bring activities to keep busy for approximately 2-4 hrs. Given the long duration, you must remain in the exam room. Your child will benefit from your attention, so consider arranging alternative care for siblings.



What medication/foods should I bring?

- For an oral challenge to a drug, your physician will let you know if you need a prescription and/or will need to bring the medication on the day of the challenge.
- **For an oral challenge to a food, you are responsible for bringing at least a serving size to be challenged. Please bring several different options to the appointment. We are only able to challenge one type of food per challenge. Please bring the original packaging for the nutritional information and ingredients. Read carefully to avoid inclusion or cross-contamination of other foods to which you/your child may be allergic.**

FOOD TO BE CHALLENGED	EXAMPLES
Baked cow's milk	*Follow <u>one</u> of the recipes (in a separate PDF)
Baked egg	*Follow recipe (in separate PDF)
cow's milk	<ul style="list-style-type: none"> • At least 8 oz of milk, yogurt – Plain or flavored • Small carton of ice cream
crab or shrimp	<ul style="list-style-type: none"> • At least 1-2 oz cooked food (prepared at home)
egg	<ul style="list-style-type: none"> • 2 scrambled eggs (1 egg/container) • 2 slices of French toast (1 egg/slice)
fish	<ul style="list-style-type: none"> • At least 1-2 oz fish prepared at home (not canned)
peanuts	<ul style="list-style-type: none"> • Peanut butter (at least 2 tablespoons) • Reese's Pieces: 84 pieces (two 6 oz bags) • Reese's Peanut Butter Cups: 4 Cups (not minis) • Peanuts: at least 30 peanuts • Bamba: two 3.5 oz packages
sesame	<ul style="list-style-type: none"> • Tahini (at least 1 oz) • <i>Sesame Snaps</i> (at least 8 candies)
soy	<ul style="list-style-type: none"> • Soy butter (at least 2 tablespoons) • At least 8 oz of soy milk, yogurt
sunflower seed	<ul style="list-style-type: none"> • Sunflower seeds (about 1/3 cup) • Sunflower seed butter (at least 2 tablespoons)
tree nut	<ul style="list-style-type: none"> • At least 2 oz of roasted nuts • Nut butter: store-bought (i.e., Zinke Orchard almond butter) or homemade using a food processor & may add vegetable oil, sugar/honey to taste
wheat	<ul style="list-style-type: none"> • Wheat cereal, crackers, bread (at least 2-3 slices)